

# Searching for AGNs among unidentified *INTEGRAL* sources

E. Maiorano<sup>1\*</sup>, R. Landi<sup>1</sup>, J.B. Stephen<sup>1</sup>, L. Bassani<sup>1</sup>, N. Masetti<sup>1</sup>, P. Parisi<sup>1</sup>,  
E. Palazzi<sup>1</sup>, P. Parma<sup>2</sup>, A.J. Bird<sup>3</sup>, A. Bazzano<sup>4</sup>, P. Ubertini<sup>4</sup>, E. Jiménez-Bailón<sup>5</sup>,  
V. Chavushyan<sup>6</sup>, G. Galaz<sup>7</sup>, D. Minniti<sup>7,8</sup>, L. Morelli<sup>9</sup>

<sup>1</sup> *IASF/INAF Bologna, via Piero Gobetti 101, 40129 Bologna, Italy*

<sup>2</sup> *IRA/INAF Bologna, via Piero Gobetti 101, 40129 Bologna, Italy*

<sup>3</sup> *School of Physics and Astronomy, University of Southampton, SO17 1BJ, UK*

<sup>4</sup> *IASF/INAF Roma, via Fosso del Cavaliere 100, 00133 Rome, Italy*

<sup>5</sup> *Instituto de Astronomía, Universidad Nacional Autónoma de México, Apartado Postal 70-264, 04510 México D.F., México*

<sup>6</sup> *Instituto Nacional de Astrofísica, Óptica y Electrónica, Apartado Postal 51-216, 72000 Puebla, México*

<sup>7</sup> *Departamento de Astronomía y Astrofísica, Pontificia Universidad Católica de Chile, Casilla 306, Santiago 22, Chile*

<sup>8</sup> *Specola Vaticana, V-00120 Città del Vaticano*

<sup>9</sup> *Dipartimento di Astronomia, Università di Padova, Vicolo dell'Osservatorio 3, I-35122 Padua, Italy*

## ABSTRACT

We report on a new method to identify Active Galactic Nuclei (AGN) among unidentified *INTEGRAL* sources. This method consists of cross-correlating unidentified sources listed in the fourth IBIS Survey Catalogue first with infrared and then with radio catalogues and a posteriori verifying, by means of X-ray and optical follow up observations, the likelihood of these associations. In order to test this method, a sample of 8 sources has been extracted from the fourth IBIS Catalogue. For 7 sources of the sample we obtained an identification, whereas the last one (IGR J03103+5706) has insufficient information for a clear classification and deserves more in-depth study. We identified three objects (IGR J08190–3835, IGR J17520–6018, IGR J21441+4640) as AGNs and suggest that three more (IGR J00556+7708, IGR J17219–1509, IGR J21268+6203) are likely active galaxies on the basis of their radio spectra, near-infrared photometry and location above the Galaxy plane. One source (IGR J05583–1257) has been classified as a starburst galaxy but it might have been spuriously associated with the *INTEGRAL* detection.

**Key words:** Catalogues, Surveys, Gamma-Rays: Observations

## 1 INTRODUCTION


A key strategic objective of the *INTEGRAL* mission (Winkler et al. 2003) is a survey of the sky at high energies ( $> 20$  keV), the domain where fundamental changes from primarily thermal to non-thermal sources/phenomena are expected, where the effects of absorption are drastically reduced and where most of the extreme astrophysical behaviour are taking place. To survey the high energy sky, *INTEGRAL* makes use of the unique imaging capability of the IBIS instrument (Ubertini et al. 2003), which allows the detection of sources at the mCrab flux level with an angular resolution of  $12'$  and a point source location accuracy of typically  $1\text{--}3'$  within a large ( $29 \times 29$  degrees) field of view.

So far, several surveys produced from data collected by IBIS have been reported in the literature, the most complete

being that of Bird et al. (2010), which lists more than 700 sources of a diverse nature (Galactic and extra-galactic) and class. However, a large fraction ( $\sim 30\%$ ) of these new *INTEGRAL* sources has no obvious counterpart in other wavebands and cannot be firmly classified; their classification is a primary objective of the survey work but it is made very difficult by the large IBIS error circles. Improved arcsecond-sized localization is therefore necessary to pinpoint the optical counterpart and through spectroscopic observations assess its nature/class (Masetti et al. 2010). For source identification one relies mostly on X-ray observations, but data in other wavebands can be used as well for counterpart search, in particular in those cases where the *INTEGRAL* unidentified source may be associated with an Active Galactic Nucleus (AGN).

Within this framework, in the present paper we present a new method for AGN identification in the 4th IBIS catalogue, which relies on cross-correlating unidentified sources

\* E-mail: maiorano@iasfbo.inaf.it



**Figure 1.** NVSS image cut-outs for four sources in our sample with overimposed IBIS error circle and 2MASX source positions. In all images, the north is up and east to the left. The scale is reported at the bottom right corner.

first with infrared (IR) and then with radio catalogues. The method is a posteriori verified by means of X-ray and optical follow up observations of some of the sources discussed in the paper, which allow exploring their nature.

In the first step we use a set of extended IR objects (2MASS Extended Catalogue, Skrutskie et al. 2006) 97% of which are associated with galaxies; we note that this is one of the most complete lists of galaxies available as it covers also the Galactic plane. Then we search for radio emission from these galaxies in order to identify likely AGN. This second step is justified by the fact that almost all AGN detected so far by *INTEGRAL* have a radio counterpart, which is not necessary radio-loud but can emit at a few mJy level.

An object which is extragalactic in nature and radio emitting, could also be a starburst galaxy but, in this case, we would not expect a bright X-ray luminosity as typically seen above 20 keV with currently operating hard X-ray detectors; indeed, so far, no starburst galaxy has been detected by IBIS in the 20–100 keV energy band.


Hence we use the IR, or better the information on source extension provided by the 2MASS extended catalogue, to pinpoint galaxies and then the information available from radio and hard X-ray emissions to look for AGN. We emphasize that many unidentified objects in the IBIS survey are on the Galactic plane, so that identification of these sources as AGN is not straightforward. Our method however provides a way to overcome problems related to the identification of active galaxies in the “zone of avoidance”. Our choice of IR and radio catalogues is not intended to be a way to select special types of AGN, but rather a simple way to

find galaxies among galactic and extragalactic sources and to select, among them, AGN associated with hard X-ray selected objects. X-ray follow up observations are instead used to provide confirmation of the proposed IR/radio and hard X-ray association while optical spectroscopy is performed to test the source AGN nature and class.

The paper is organized as follows: Sect. 2 describes the sample selection criteria and gives an overview of the extracted sample; the multiwavelength observations and analyses are reported in Sect. 3, while results and discussion are shown in Sect. 4. Finally, conclusions are presented in Sect. 5. The present work supersedes the analysis carried out in Maiorano et al. (2010), in which preliminary results for only three sources of the sample were presented.

## 2 SAMPLE SELECTION

Historically, AGNs were discovered with radio observations, i.e. the radio selection is often a way to recognize active galaxies, except at lower luminosities where star-formation in galaxies can produce radio emission. Therefore, for bright objects, a mere detection in radio provides support for the presence of an active galaxy. Sample contamination from Galactic sources may however come from pulsars, microquasars and Cataclismic Variables (CVs). In many cases, association with a galaxy via cross-correlation with galaxy catalogues can help in selecting only extragalactic objects and, by means of the radio detection, pinpointing those sources that are likely AGNs.



**Figure 2.** NVSS or SUMSS image cut-outs for four sources in our sample with overimposed IBIS error circle and 2MASX source positions. In all images, the north is up and east to the left. The scale is reported at the bottom right corner.

**Table 1.** IBIS, 2MASX and radio positions (and corresponding errors) for each source in the sample. The radio positions come from the analysis carried out in this work (Subsect. 2.1).

Source	IBIS Position RA (J2000) Dec (J2000)	IBIS error circle (arcmin)	2MASX Object	2MASX Position RA (J2000) Dec (J2000)	Radio Object	Radio Position RA (J2000)(err) Dec (J2000)(err)
IGR J00556+7708	00:55:34.8 +77:08:24	4.9'	2MASX J00570148+7708505	00:57:01.482 +77:08:50.50	NVSS J005700+770911	00:57:00.016 (0.181s) +77:09:11.81 (0.62'')
IGR J03103+5706	03:10:16.8 +57:06:07.2	4.9'	2MASX J03095498+5707023	03:09:54.987 +57:07:02.34	NVSS J030954+570704	03:09:54.955 (0.128s) +57:07:04.29 (1.04'')
IGR J05583-1257	05:58:17.04 -12:57:43.2	4.8'	2MASX J05580231-1255477 (LCSB 02890)	05:58:02.313 -12:55:47.78	NVSS J055802-125545	05:58:02.416 (0.172s) -12:55:45.54 (3.30'')
IGR J08190-3835	08:19:02.16 -38:34:58.8	3.5'	2MASX J08191136-3833104	08:19:11.365 -38:33:10.46	NVSS J081910-383307	08:19:10.929 (0.219s) -38:33:06.78 (3.00'')
IGR J17219-1509	17:21:55.68 -15:09:39.6	4.2'	2MASX J17215337-1505384	17:21:53.379 -15:05:38.49	NVSS J172153-150532	17:21:53.273 (0.155s) -15:05:32.02 (3.70'')
IGR J17520-6018	17:52:02.16 -60:18:18	5.0'	2MASX J17515581-6019430	17:51:55.818 -60:19:43.08	SUMSS J175155-601943	17:51:55.585 (0.142s) -60:19:43.87 (1.30'')
IGR J21268+6203	21:26:46.08 +62:03:43.2	4.4'	2MASX J21262644+6204410	21:26:26.440 +62:04:41.03	NVSS J212628+620457	21:26:28.707 (0.027s) +62:04:57.58 (0.19'')
IGR J21441+4640	21:44:04.08 +46:40:51.6	4.9'	2MASX J21441345+4637169 (UGC 11806)	21:44:13.455 +46:37:16.97	NVSS J214413+463718	21:44:13.419 (0.092s) +46:37:17.79 (0.98'')
			2MASX J21435408+4637048 (UGC 11802)	21:43:54.082 +46:37:04.84	NVSS J214354+463705	21:43:54.055 (0.074s) +46:37:04.99 (0.66'')

So, while a mere radio detection does not imply an identification with an AGN, its combination with high energy X/gamma-ray emission, together with the association with a galaxy, strongly argues in favour of the identification of an unclassified *INTEGRAL* source with an active galaxy. Following this reasoning we have cross-correlated our set of

unidentified *INTEGRAL* sources in the 4th IBIS catalogue (Bird et al. 2010) with IR/radio catalogues, in order to extract a small sample of objects likely associated with an AGN.

For the IR bands we have used the Two Micron All Sky Survey Extended (2MASX) Source Catalog (Skrutskie et al.

2006) which is a powerful tool to identify, within the sample of unidentified *INTEGRAL* sources, those possibly associated with galaxies. For this survey, the entire sky was uniformly scanned in three near-infrared (NIR) bands (J,H,K) to detect and characterise sources brighter than about 1 mJy in each band, with a signal-to-noise ratio greater than 10 and which are resolved and extended beyond the Two Micron All Sky Survey (2MASS) beam/point spread function. The absolute astrometric accuracy of the 2MASX catalogue is better than one arcsec. The extended source catalogue consists of 1,647,599 objects, 97% of which are galaxies while the remaining  $\sim 3\%$  is made of sources in the Milky Way (mostly double and triple stars, HII regions, planetary and reflection nebulae), which are not expected to emit at high energies. Therefore, the presence of a 2MASX object inside the IBIS error circle suggests an association with a galaxy.

As radio catalogues we have used the NRAO VLA Sky Survey (NVSS, Condon et al. 1998) and the Sydney University Molonglo Sky Survey (SUMSS, Mauch et al. 2003) which are particularly well suited for finding counterparts of unidentified *INTEGRAL* sources: they are similar in sensitivity and spatial resolution and together they cover the whole sky. The NVSS catalogue covers the sky north of the J2000.0 Declination of  $-40$  degrees (82% of the celestial sphere) at 1.4 GHz (20 cm). The catalogue consists of almost 2 million discrete sources stronger than a flux density of about 2.5 mJy. The NVSS images have 45 arcsecond FWHM angular resolution and nearly uniform sensitivity. The *rms* uncertainties in right ascension and declination vary from about  $1''$  for the 400,000 sources stronger than 15 mJy to  $7''$  at the survey limit. The SUMSS catalogue covers instead the sky south of the J2000.0 Declination of  $-30$  degrees ( $\sim 20\%$  of the celestial sphere) and is carried out at 843 MHz (36 cm). The survey consists of  $4.3^\circ \times 4.3^\circ$  mosaic images with a resolution of  $45 \times 45$  cosec $|\delta|$  arcsec $^2$ , and a *rms* noise level of 1-2 mJy/beam. Positions in the catalogue are accurate to within  $1-2''$  for sources with peak brightness  $> 20$  mJy/beam, and are always better than  $10'$ . The internal flux density scale is accurate to within 3%. The radio detection of a 2MASX emitting galaxy strongly indicates that the source is an AGN if also detected above 10 keV.

To perform the correlation, we used the standard statistical technique which has been employed very successfully in other cases (Stephen et al. 2005, 2006, 2010). This consists of simply calculating the number of *INTEGRAL* sources for which at least one 2MASX counterpart was within a specified distance, out to a distance where all *INTEGRAL* sources had at least one NIR counterpart. To have a control group we created a list of fake “anti-*INTEGRAL*” sources. For every object in the *INTEGRAL* list, we made a corresponding source in the fake list with coordinates mirrored in Galactic longitude and latitude (this mirroring was chosen due to the strong Galactic component evident in the *INTEGRAL* distribution), and the same correlation algorithm was then applied between this list and the 2MASX catalogue. Subtracting from the number of correlations in the true list those obtained in the false sample, it is possible to estimate the number of true associations. We see that the radius at which the first correlations between the “anti-*INTEGRAL*” sources sample and 2MASX catalogue appear is about 6 arcmin. This is comparable with the size

of the IBIS error circle radius, therefore we expect at most 1 spurious correlation within the lot of selected sources (see below). The sample of associations extracted in this way, i.e. a list of objects likely associated with galaxies, was then cross-correlated to the radio catalogues following the same method.

By means of this sequence of cross-correlations we extracted a final sample of 8 objects which are seen in all 3 wavebands (hard X-rays, NIR and radio); all 8 can be considered as AGN candidates because they are classified as galaxies in the 2MASX and are detected both in radio and hard X-rays. Note that in one case we have two radio/NIR objects associated with a unique *INTEGRAL* source (IGR J21441+4640); both are detected in the 2MASX and radio catalogues and so are equally possible counterparts of the *INTEGRAL* source and as such will be considered in the following sections. In NASA/IPAC Extragalactic Database (NED) these two objects form a galaxy pair.

## 2.1 Overview of the extracted sample

All objects in the sample are reported in Table 1; for each source we list the *INTEGRAL* name, the IBIS position with relative error, the IR (2MASX) and radio (NVSS or SUMSS) positions and associated errors of the putative counterparts. In our sample, 7 sources have a counterpart in the NVSS catalogue while only one has an association in the SUMSS.

The 2MASX catalogue provides J, H, K magnitudes for all sources in the sample which are reported in Table 2; note that all objects listed in Table 2 are classified as galaxies in NED. In Table 2 we also show the NIR colour indices J–H and H–K; these can be used as a tool to confirm the AGN nature of our sources. Indeed, all objects but one (2MASX J03095498+5707023) have NIR colours compatible with those of nearby active galaxies (see Fig. 1 of Kouzuma & Yamaoka, 2010). Despite the uncertainties introduced by this method (i.e. it is not obvious from the work of Kouzuma & Yamaoka [2010], which is the efficiency of finding an AGN via IR photometry), it is reassuring that almost all of our objects are compatible with NIR AGN colour indices.

Hard X-ray information concerning the sources of our sample is reported in Bird et al. (2010). We note here that all but two objects are classified as variable in the 4th IBIS catalogue since they are detected at the level of Revolution, Revolution Sequence or through the bursticity analysis; the only exceptions are IGR J05583–1257 and IGR J08190–3835 which are reported as persistent objects in the IBIS survey. All but 3 objects (IGR J03103+5706, IGR J08190–3835, IGR J21441+4640) are at Galactic latitude  $> 10^\circ$  an additional argument in favour of their extragalactic nature.

In order to provide radio images and fluxes, we have used the standard procedure employed within the software package AIPS<sup>1</sup>, release 31DEC10 (Astronomical Image Processing System). NVSS/SUMSS fluxes extracted from the radio maps are reported in Table 3.

Figures 1 and 2 show the collection of NVSS/SUMSS image cut-outs for all of our sources with overlaid IBIS error circle and 2MASX source positions. For two sources (IGR J00556+7708, IGR J21268+6203) in our sample the

<sup>1</sup> <http://www.aips.nrao.edu/>

radio and NIR positions are significantly different from each other, being separated by  $22''$  and  $24''$ , respectively. In both cases we searched among the optical and NIR catalogues selected in the Vizier Catalogues Selection Page<sup>2</sup> (USNO-B1, Monet et al. 2003, USNO-A2.0<sup>3</sup>, 2MASS), but we could not find any optical or NIR counterparts within the radio error circle; furthermore, we note that the 2MASX galaxies lie in each case within the edge of the radio contours and are obviously extended in the NIR band (semimajor axis of  $16.6''$  and  $13.7''$ , respectively). Indeed, in both cases the 2MASX and radio objects are associated with each other in NED on the basis of a sophisticated cross-identification analysis which takes into account not only the positional uncertainty but also the source extension (Mazzarella et al. 2001). Taking all these evidences into account, we decide to keep both *INTEGRAL* sources in the sample also in view of the fact that either the NIR or the radio object (or in the best case both) are likely counterparts to the high energy emitter.

By looking for further radio information in the data archives of different radio telescopes and also in the literature, we found that some sources in the sample have been observed at more than one radio frequency. In these few cases we have calculated the radio spectral index using the available data points and the usual relation  $F_\nu \propto \nu^{-\alpha}$  (see Sect. 3). Furthermore, when the redshift is available, we have calculated the radio power at 1.4 GHz; all these values are reported in Table 3. Note that all objects in Table 3 have compact morphology in radio.

### 3 FOLLOW UP OBSERVATIONS

In order to test the validity of our method as well as to confirm the AGN nature of our objects, we have obtained a set of X-ray and optical follow up observations. The X-ray observations, carried out with Swift/XRT<sup>4</sup>, were useful to confirm the association between the NIR/radio source and the IBIS hard X-ray emission and hence to pinpoint the optical counterpart; while optical spectra of this counterpart allowed us to assess the source AGN nature and class.

For four objects in our sample (see Table 4), we have X-ray observations acquired with the X-ray Telescope (XRT, 0.2–10 keV, Burrows et al. 2005) on board the *Swift* satellite (Gehrels et al. 2004). XRT data reduction was performed using the XRTDAS standard data pipeline package (XRT-PIPELINE v. 0.12.4), in order to produce screened event files. All data were extracted only in the Photon Counting (PC) mode (Hill et al. 2004), adopting the standard grade filtering (0–12 for PC) according to the XRT nomenclature.

Events for spectral analysis were extracted within a circular region of radius  $20''$ , centered on the source position, which encloses about 90% of the PSF at 1.5 keV (see Moretti et al. 2004).

The background was taken from various source-free regions close to the X-ray source of interest, using circular regions with different radii in order to ensure an evenly sampled background. In all cases, the spectra were extracted from the corresponding event files using the XSELECT software and binned using GRPPHA in an appropriate way, so that the  $\chi^2$  statistic could be applied. We used version v.011 of the response matrices and create individual ancillary response files *arf* using XRTMKARF v. 0.5.6.

The data have been fitted using an absorbed powerlaw model; due to the poor statistical quality of the X-ray data, we have fixed the photon index to 1.8 in order to evaluate the presence of absorption and the 2–10 keV flux.

For four sources in the sample (see Table 5) optical spectroscopy of the proposed counterparts was obtained from data collected at the 1.5-m telescope of the Cerro Tololo Interamerican Observatory (CTIO), Chile, at the 2.1-m telescope of the Observatorio Astronómico Nacional in San Pedro Mártir (SPM), México, and from the Six-degree Field Galaxy Survey<sup>5</sup> (6dFGS) archive (Jones et al. 2004) containing spectra acquired with the 4-m Anglo-Australian Telescope (AAT) in Siding Spring, Australia. Table 5 reports the log of these observations.

The spectroscopic data acquired at these telescopes were optimally extracted (Horne 1986) and reduced following standard procedures using IRAF<sup>6</sup>. Calibration frames (flat fields and bias) were taken on the day preceeding or following the observing night. The wavelength calibration was performed using lamp data acquired soon after each on-target spectroscopic acquisition; the uncertainty in this calibration was  $\sim 0.5$  Å in all cases, according to our checks made using the positions of background night sky lines. Flux calibration was performed using catalogued spectrophotometric standards.

As mentioned above an additional spectrum, for 2MASX J05580231–1255477, was retrieved from the 6dFGS archive. Since this archive provides spectra that are not flux-calibrated, we used the optical photometric information in Jones et al. (2005) to calibrate the 6dFGS spectrum presented in this work.

The flux calibration was obtained by normalizing the count spectrum and by multiplying it by a cubic spline constructed with the fluxes extracted from the *BRI* optical magnitudes available for the source using the conversion formulae of Fukugita et al. (1995).

The results of these X-ray and optical follow up observations are presented in Table 4 and Table 6.

### 4 RESULTS

In the following, the results of all available archival information gathered on each individual source are discussed together with the X-ray and optical data when available.

As for the X-ray data, we note that in 3 cases (IGR

<sup>2</sup> <http://vizier.u-strasbg.fr/viz-bin/VizieR>

<sup>3</sup> <http://www.nofs.navy.mil/projects/pmm/USNOSA2doc.html>

<sup>4</sup> The Swift/XRT observations were performed in the context of the approved follow-up program in collaboration with the Swift team.

<sup>5</sup> <http://www.aao.gov.au/local/www/6df/>

<sup>6</sup> IRAF is the Image Reduction and Analysis Facility made available to the astronomical community by the National Optical Astronomy Observatories, which are operated by AURA, Inc., under contract with the US National Science Foundation. It is available at <http://iraf.noao.edu>

**Table 2.** IR magnitudes as quoted in the 2MASS extended catalogue for all objects in the sample. In the case of IGR J21441+4640, we report the IR magnitudes and colours of both galaxies in the pair individually.

Source	J-mag	H-mag	K-mag	J-H	H-K
2MASX J00570148+7708505	14.922±0.222	14.291±0.333	13.270±0.173	0.63	1.02
2MASX J03095498+5707023	15.497±0.301	13.831±0.143	12.772±0.096	1.67	1.06
2MASX J05580231-1255477	13.192±0.080	12.608±0.119	12.111±0.133	0.58	0.50
2MASX J08191136-3833104	12.529±0.058	11.303±0.037	10.818±0.049	1.23	0.49
2MASX J17215337-1505384	15.563±0.237	>14.520	13.985±0.181	1.04	0.53
2MASX J17515581-6019430	14.525±0.133	13.564±0.126	13.408±0.166	0.96	0.15
2MASX J21262644+6204410	14.777±0.185	14.172±0.234	13.314±0.174	0.6	0.86
2MASX J21441345+4637169 (UGC 11806)	12.106±0.051	11.530±0.062	11.312±0.074	0.57	0.22
2MASX J21435408+4637048 (UGC 11802)	11.814±0.055	11.088±0.060	10.743±0.064	0.73	0.34

**Table 3.** Radio information of all objects in the sample. NVSS 1.4 GHz (20 cm) and SUMSS 843 MHz (36 cm) fluxes are from this work. Other flux values are from the Westerbork Northern Sky Survey (WENSS) at 92 cm and the Parkes MIT-NRAO (PMN) Surveys at 6 cm. All the redshifts reported in the Table are from NED, except for NVSS J081910-383307 (2MASX J08191136-3833104) that is derived from the spectroscopic data analysis carried out in this work (Sect. 3).

Source	Redshift	325 MHz 92 cm (mJy)	843 MHz ~ 36 cm (mJy)	1400 MHz 20 cm (mJy)	4850 MHz ~ 6 cm (mJy)	Spectral Index $\alpha_{325\text{MHz}}^{1.4\text{GHz}}$ ( $F_\nu \propto \nu^{-\alpha}$ )	Radio Power at 1.4 GHz ( $\text{WHz}^{-1}$ )
NVSS J005700+770911		51±2.1		16.0±1.2		0.79 <sup>+0.09</sup> <sub>-0.08</sub>	
NVSS J030954+570704		21±3.8		10.2±1.0		0.50 <sup>+0.18</sup> <sub>-0.20</sub>	
NVSS J055802-125545	0.003042			4.7±0.8			9.31×10 <sup>19</sup>
NVSS J081910-383307	0.009			2.7±0.7			4.71×10 <sup>20</sup>
NVSS J172153-150532				5.2±1.0			
SUMSS J175155-601943			25.2±2.2				
NVSS J212628+620457		29±4.8		45.1±1.4	19±4	-0.30 <sup>+0.13</sup> <sub>-0.16</sub>	
NVSS J214413+463718 (UGC 11806)	0.011081	23±4		11.0±1.1		0.51 <sup>+0.17</sup> <sub>-0.19</sub>	2.92×10 <sup>21</sup>
NVSS J214354+463705 (UGC 11802)	0.010517	40±4		22.9±1.2		0.38 <sup>+0.09</sup> <sub>-0.10</sub>	5.47×10 <sup>21</sup>

**Table 4.** Spectral parameters derived from the X-ray data analysis of four sources observed by Swift/XRT. The values of the Galactic column densities are from Kalberla et al. (2005).

Source	Exposure (s)	Count rate (counts/s)	$\Gamma$ Photon index	$N_{\text{H}}(\text{Gal})$ ( $\text{cm}^{-2}$ )	$N_{\text{H}}(\text{intr})$ ( $\text{cm}^{-2}$ )	Flux(2-10 keV) ( $\text{erg cm}^{-2}\text{s}^{-1}$ )
IGR J05583-1257 (LCSB 02890)	4623	<0.7×10 <sup>-3</sup>	1.8 (fixed)	1.63×10 <sup>21</sup>		<5.8×10 <sup>-14</sup>
IGR J08190-3835	10310	(18.6±1.3)×10 <sup>-3</sup>	1.8 (fixed)	9.6×10 <sup>21</sup>	13.6×10 <sup>22</sup>	1.49×10 <sup>-12</sup>
IGR J17520-6018	11883	(18.2±1.2)×10 <sup>-3</sup>	1.8 (fixed)	7.0×10 <sup>20</sup>	13×10 <sup>22</sup>	2.55×10 <sup>-12</sup>
IGR J21441+4640 (UGC 11806)	2755	(3.3±1.2)×10 <sup>-3</sup>	1.8 (fixed)	2.57×10 <sup>21</sup>		1.2×10 <sup>-13</sup>
(UGC 11802)		<1.2×10 <sup>-3</sup>	1.8 (fixed)	2.8×10 <sup>21</sup>		<8.3×10 <sup>-14</sup>

**Table 5.** Log of the spectroscopic observations presented in this paper.

Object	Date	Telescope + instrument	Exp. start time (UT)	Disp. (Å/pix)	Exposure time (s)
LCSB L0289O	13 Jan 2005	AAT+6dF	12:15	1.6	1200+600
USNO-A2.0 0450_06519994	18 Jan 2010	CTIO 1.5m + RC Spec.	03:54	5.7	2×1200
UGC 11802	15 Sep 2009	SPM 2.1m + B&C Spec.	04:37	4.0	2×1800
UGC 11806	15 Sep 2009	SPM 2.1m + B&C Spec.	05:59	4.0	2×1800

J08190-3835, IGR J17520-6018 and IGR J21441+4640) the X-ray measurements confirm the proposed counterpart; in the case of the galaxy pair only one of the two objects, UGC 11806, was detected while an upper limit was set on the other, UGC 11802. Only in one case (IGR J05583-1257) no X-ray source is detected within the *INTEGRAL* error circle (see Section on this source) and an upper limit is reported for the proposed counterpart.

For the optical spectra categorization, as they all refer to extragalactic sources (see below), we used the criteria of Veilleux & Osterbrock (1987) and the line ratio diagnostics of both Ho et al. (1993, 1997) and Kauffmann et al. (2003) which are generally used for emission-line AGN classification. The spectra of the galaxies shown here were not corrected for starlight contamination (see, e.g., Ho et al. 1993, 1997) because of the limited S/N and spectral resolution. In this case, we do not expect this to affect any of our main results and conclusions.

Of the four objects for which we have obtained optical spectra, one was found to be a type 2 AGN (IGR J08190-3835), two (i.e. those belonging to the galaxy pair) were classified as LINERs and one shows the features typical of a starburst galaxy (IGR J05583-1257).

In the following, we consider a cosmology with  $H_0 = 71$  km s<sup>-1</sup> Mpc<sup>-1</sup>,  $\Omega_\Lambda = 0.73$ , and  $\Omega_m = 0.27$ ; the luminosity distances of the extragalactic objects reported in this paper were computed for these parameters using the Cosmology Calculator of Wright (2006).

#### IGR J00556+7708

This source, located at high Galactic latitude, is detected by IBIS in a Revolution Sequence suggesting that it might be variable on long timescales. As said above, in this case the radio and NIR positions are significantly different and the objects are located on the left edge of the IBIS error box (Fig. 1, top-left panel). The radio object is detected at 20 and 92 cm and has a steep spectrum with index  $\alpha=0.79$ , typical of a radio loud AGN; no optical counterpart is found within the radio source positional uncertainty. The NIR source is also listed in the USNO-A2.0 catalogue<sup>7</sup> with name USNO-A2.0 1650\_00208794 and optical magnitudes  $R \sim 17.4$  and  $B \sim 19.5$ ; the NIR colour indices are typical of an active galaxy (Kouzuma & Yamaoka, 2010).

Overall, we conclude that the radio and NIR objects are both likely AGN and, as such, are both possible counterparts of the IBIS detection.

#### IGR J03103+5706

This object is also likely variable as it was detected by IBIS during just one revolution (3 days). The radio/NIR counterpart suggested in this work is located within the IBIS error circle (Fig. 1, top-right panel). It is detected in radio at 20 and 92 cm and has  $\alpha=0.50$ , which is characteristic of a radio flat spectrum galaxy. The source has no optical counterpart in USNO-B1.0 (Monet et al. 2003), suggestive of a highly reddened/absorbed object. Its NIR photometry, as well as its location on the Galactic plane, are not typical of an AGN, so the nature of this object remains dubious.

#### IGR J05583-1257

This is one of the two persistent sources in the sample; the radio/NIR source is identified in NED with LCSB L0289O, a low brightness galaxy at  $z = 0.003$ . This source is fairly bright in the far IR being detected by IRAS at 60  $\mu$ m with a flux of 0.6 Jy (IRAS, 1988). The available radio data (Fig. 1, bottom left panel) for this source only consist of a flux at 20 cm, thus no information on the spectral index can be obtained. The 6dFGS optical spectrum (Fig. 4, first panel from top; see also DSS-II-red image in Fig. 3, left panel) shows several permitted and forbidden narrow emission lines at the above redshift; their ratios (see also Table 6) indicate that this is a starburst galaxy with no noticeable nuclear activity.

There is no X-ray detection within the 90% IBIS error box, but at the border of the 99% positional uncertainty we find a detection at low significance level,  $2.6\sigma$ . This source is located at RA(J2000) = 05h 57m 59.9s, Dec(J2000) =  $-12^\circ 51' 33.0''$  ( $6''$  uncertainty). The X-ray flux is  $7.3 \times 10^{-14}$  erg cm<sup>-2</sup> s<sup>-1</sup>, with photon index fixed to 1.8. At this position no optical or IR or radio source is found. On the other hand, we do not find any X-ray emission at the location of LCSB 0289O. The 2–10 keV flux upper limit provides an indication on the source luminosity which is  $< 10^{39}$  erg s<sup>-1</sup>, i.e. quite low for an AGN, unless the source is extremely variable (but this is not evident in the IBIS data), or extremely absorbed (thus masking a Compton thick AGN in a starburst galaxy).


The persistent nature of the source, the stringent upper limit in X-rays and the optical spectrum suggest that this is probably not the counterpart of the IBIS source, which is either a spurious detection or has a different association than LCSB 0289O.

#### IGR J08190-3835

This is the second persistent source in our sample. Within the IBIS uncertainty, XRT detects only one source with a statistical significance of  $5.8\sigma$  in the energy range 0.3–

<sup>7</sup> <http://www.nofs.navy.mil/projects/pmm/USNOSA2doc.html>





**Figure 3.** Optical images of the fields of 3 of the *INTEGRAL* hard X-ray sources selected in this paper for optical spectroscopic follow-up: IGR J05583–1257 (left panel), IGR J08190–3835 (central panel) and IGR J21441+4640 (right panel). The proposed optical counterparts are indicated with tick marks (see text). Field sizes are  $10' \times 10'$  and are extracted from the DSS-II-Red survey. In all cases, north is up and east to the left.

10 keV. This object is positionally coincident with both 2MASX and NVSS sources (Fig. 1, bottom-right panel). The radio analysis provides a flux of  $\sim 3$  mJy at 20 cm. The source is fairly bright at NIR frequencies (see Table 2) and it is also detected in the optical band, where it is listed in the USNO-A2.0 catalogue with name USNO-A2.0 0450\_06519994 (see Table 5 and Table 6) and magnitudes  $R \sim 14.6$  and  $B \sim 16.8$ . From a comparison with the NIR magnitudes in Table 2, the  $R-K$  colour is therefore  $\sim 4$ , which suggests a red or obscured galaxy (Fig. 3, central panel and Table 6).

The X-ray data analysis is described by an absorbed power law with photon index fixed to 1.8 and an observed 2–10 keV flux of  $1.5 \times 10^{-12}$  erg cm $^{-2}$  s $^{-1}$ ; the intrinsic column density is  $N_H(intr) \sim 1.4 \times 10^{23}$  cm $^{-2}$ , which exceeds the Galactic value of  $9.6 \times 10^{21}$  cm $^{-2}$  (Kalberla et al. 2005). Optical spectroscopy (Fig. 4, second panel from top) shows that the source displays  $H_\alpha$ , [N II] and [S II] narrow emission lines at redshift  $z = 0.009 \pm 0.001$  superimposed on a very reddened continuum. The flux ratios among these emission features suggest that the object is a Type 2 AGN.

Overall, we conclude that the proposed 2MASX/NVSS/XRT source is the actual counterpart of the IBIS-detected object and it is a Narrow Line (obscured) AGN.

#### IGR J17219–1509

The radio/NIR source is located at the edge of the IBIS error circle (Fig. 2, top-left panel); it is very weak in radio, being close to the flux limit of the NVSS map ( $\sim 5$  mJy). Very little is known about this source except that it is strongly variable in the IBIS waveband, being detected only in one revolution (R403, i.e. for a few days) and showing a large bursticity factor of 4 (see Bird et al. 2010 for details). The source has an optical magnitude of 18.2 in  $R$ ; the NIR colours, as well as its location above the Galaxy plane, are compatible with the source being a nearby AGN (Kouzumá & Yamaoka, 2010).

We conclude that this is an extragalactic object, most likely an AGN, on the basis of the above considerations and it is a good association for the *INTEGRAL* source.

#### IGR J17520–6018

This source can also be considered as variable by Bird et

al. (2010). An X-ray source is well detected at  $13.6\sigma$  confidence level in the range 0.3–10 keV by XRT within the IBIS uncertainty and is located at  $RA(J2000) = 17^h 51^m 55.9^s$  and  $Dec(J2000) = -60^\circ 19' 44''$  ( $4''$  uncertainty). It coincides with both the 2MASS extended object and the radio source reported in the SUMSS (Fig. 2, top-right panel). The source has only one detection in radio at 36 cm. Besides being detected in the NIR, the galaxy is also listed in the USNO-B1 catalogue with optical B and R magnitudes of 15.5 and 14.7, respectively. The X-ray data analysis provides an absorbed power law spectrum with fixed photon index of 1.8 and an observed 2–10 keV flux of  $2.6 \times 10^{-12}$  erg cm $^{-2}$  s $^{-1}$ ; the intrinsic column density is  $N_H(intr) = 1.3 \times 10^{23}$  cm $^{-2}$  which exceeds the Galactic value (see Table 4). This source is also reported in the BAT 58–Month catalogue<sup>8</sup> as SWIFT J1751.8–6019, with flux  $1.7 \times 10^{-11}$  erg cm $^{-2}$  s $^{-1}$  in the range 14–195 keV.

Based on all the above information, we conclude that this galaxy is the likely counterpart of the IBIS/BAT source and, on the basis of the detected X-ray absorption, we further suggest that it is a type 2 AGN.

#### IGR J21268+6203

This IBIS source is also highly variable being reported with a high bursticity factor by Bird et al. (2010); it is also likely an extragalactic source being located above the Galactic plane. As said in Sect. 2.1, in this case the radio and NIR positions are significantly different (Fig. 2, bottom-left panel). The NIR object (Table 2) has an optical counterpart in USNO-B1.0 1520\_0331686 listed in USNO-B1.0 catalogue with magnitudes  $B \sim 20.3$ ,  $R \sim 17.5$  and  $I \sim 16.4$ ; no optical association was found for the radio object. The NIR colours are also in this case typical of a nearby AGN (Kouzumá & Yamaoka, 2010). The radio data available provide a spectral index  $\alpha = -0.3$  between 20 and 92 cm, indicative of a probable GHz-Peaked-Spectrum (GPS) radio source, i.e. a source characterised by a convex radio spectrum peaking near 1 GHz (Stanghellini 2006; Lister 2003; O’Dea 1998). GPS are compact powerful young radio galaxies that reside in gas-rich environments at the center of active galaxies.

<sup>8</sup> <http://swift.gsfc.nasa.gov/docs/swift/results/bs58mon/>



**Table 6.** Fluxes (in units of  $10^{-14}$  erg cm $^{-2}$  s $^{-1}$ ) of the main emission lines detected in the spectra of the objects reported in Fig. 4. The correction for the Galactic reddening was computed a color excess  $E(B - V) = 0.479, 1.54, 0.404$  and  $0.403$  mag for LCSB L02890, USNO-A2.0 0450\_06519994, UGC 11802 and UGC 11806, respectively (from Schlegel et al. 1998). Uncertainties and upper limits for the fluxes are reported at  $1\sigma$  and  $3\sigma$  confidence levels, respectively.


Line	Observed flux	Corrected flux
LCSB L02890 ( $z = 0.003$ )		
H $\beta$	$17.5 \pm 0.9$	$87 \pm 4$
[O III] $\lambda 5007$	$92 \pm 3$	$428 \pm 13$
[O I] $\lambda 6300$	$< 0.5$	$< 2.2$
H $\alpha$	$34.2 \pm 1.0$	$105 \pm 3$
[N II] $\lambda 6583$	$1.0 \pm 0.3$	$3.3 \pm 1.0$
[S II] $\lambda 6716$	$1.8 \pm 0.3$	$5.5 \pm 0.9$
[S II] $\lambda 6731$	$1.2 \pm 0.2$	$3.6 \pm 0.6$
USNO-A2.0 0450_06519994 ( $z = 0.009$ )		
H $\beta$	$< 0.07$	$< 9$
[O III] $\lambda 5007$	$< 0.07$	$< 9$
[O I] $\lambda 6300$	$< 0.17$	$< 7$
H $\alpha$	$0.32 \pm 0.08$	$11 \pm 3$
[N II] $\lambda 6583$	$0.12 \pm 0.03$	$4.1 \pm 1.0$
[S II]*	$0.26 \pm 0.04$	$8.3 \pm 1.2$
UGC 11802 ( $z = 0.0105$ )		
H $\beta$	$1.41 \pm 0.14$	$5.4 \pm 0.5$
[O III] $\lambda 5007$	$0.76 \pm 0.11$	$2.7 \pm 0.4$
[O I] $\lambda 6300$	$0.24 \pm 0.06$	$0.55 \pm 0.14$
H $\alpha$	$9.5 \pm 0.3$	$24.1 \pm 0.7$
[N II] $\lambda 6583$	$2.47 \pm 0.12$	$6.4 \pm 0.3$
[S II] $\lambda 6716$	$2.26 \pm 0.16$	$5.6 \pm 0.4$
[S II] $\lambda 6731$	$1.58 \pm 0.11$	$3.9 \pm 0.3$
UGC 11806 ( $z = 0.0110$ )		
H $\beta$	$1.22 \pm 0.12$	$4.8 \pm 0.5$
[O III] $\lambda 5007$	$0.76 \pm 0.08$	$2.8 \pm 0.3$
[O I] $\lambda 6300$	$0.21 \pm 0.05$	$0.70 \pm 0.18$
H $\alpha$	$7.0 \pm 0.2$	$17.7 \pm 0.5$
[N II] $\lambda 6583$	$2.88 \pm 0.14$	$7.2 \pm 0.4$
[S II] $\lambda 6716$	$1.53 \pm 0.11$	$3.7 \pm 0.3$
[S II] $\lambda 6731$	$1.24 \pm 0.09$	$3.1 \pm 0.3$

\*: The doublet is blended due to the low spectral S/N and resolution.

Thus, the observational evidence suggests that the radio and NIR objects are both likely AGN and, as such, are good candidates for an association with the *INTEGRAL* source.

#### IGR J21441+4640

This also is a strongly variable source in IBIS (Bird et al. 2010) with a bursticity larger than 4. Within the IBIS positional uncertainty of this source there are two galaxies (see Fig. 3, right panel), UGC 11802 and UGC 11806, at the same redshift ( $z=0.011$ ), which form the galaxy pair KPG559. Both galaxies are detected at radio frequencies in the NVSS catalogue (NVSS J214354+463705, NVSS J214413+463718) and are also listed in the 2MASS extended catalogue (2MASX J21435408+4637048, 2MASX



**Figure 4.** Spectra (not corrected for the intervening Galactic absorption) of the possible optical counterparts of some of the *INTEGRAL* sources presented in this paper (IGR J05583–1257, IGR J08190–3835 and IGR J21441+4640; see text). For each spectrum, the main spectral features are labeled. The symbol  $\oplus$  indicates atmospheric telluric absorption bands.

J21441345+4637169) (see Fig. 2, bottom-right panel). The radio data analysis provides a 20 cm flux for both objects, which are also detected at 92 cm and have spectral indices  $\alpha=0.38$  and  $\alpha=0.51$ , respectively; both radio power and spectral index are typical of low luminosity AGN. The optical B (R) magnitudes of the two objects are 14.5 (9.5) and 14.7 (12.9). Both are detected with AKARI in the far IR, from 90 to 160  $\mu\text{m}$  at the few Jansky flux level (Murakami et al. 2007). Within the IBIS uncertainty, XRT finds only one galaxy (UGC 11806) of the pair, with a low statistical significance of  $2.5\sigma$  in the range 0.3–10 keV. The XRT spectrum can be described by an unabsorbed power law with fixed photon index of 1.8 and an observed 2–10 keV flux of  $1.2 \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$ . The upper limit of the X-ray flux in the 2–10 keV band for the companion galaxy UGC11802 is  $8.3 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$ .

Optical spectroscopy (Fig. 4, lower panel) indicates that UGC 11806 is a narrow emission-line galaxy with flat continuum and prominent Balmer, [N II], [O III] and [S II] lines at a redshift consistent with the one found in the literature. Emission line ratios suggest that this is a transition object, that is, a LINER (Heckman 1980) with a possible contamination from an underlying starburst event. The optical spectrum of UGC 11802 (Fig. 4, third panel from top) indicates instead that this galaxy is a starburst, with no indication of AGN activity (which explains the absence of detectable X-ray emission from its nucleus).

The low significance detection in X-rays of UGC 11806 may be due to the variable nature of the source (Bird et al. 2010) rather than to the high absorption, which is not readily apparent from the optical spectrum. To this aim we can infer the reddening local to the source by considering an intrinsic  $H_\alpha/H_\beta$  line ratio of 2.86 (Osterbrock 1989). The corresponding color excess, obtained by comparing the intrinsic line ratio with the measured one by applying the Galactic extinction law of Cardelli et al. (1989), is  $E(B - V) = 0.25$  mag. This, using the formula of Predehl & Schmitt (1995), corresponds to a hydrogen column density  $N_H = 1.4 \times 10^{21} \text{ cm}^{-2}$  local to the AGN.

Overall we conclude that UGC 11806 is the counterpart of the IBIS source; it is probably a low luminosity and highly variable AGN of LINER type.

## 5 CONCLUSION

The basic idea of this work is to propose a way whereby AGN can be easily found out among a set of unidentified objects detected in hard X-ray surveys. The method, which consists of two consecutive steps of cross-correlations between hard X-ray objects and IR/radio catalogues, is tested here for the unidentified sources contained in the 4th IBIS survey catalogue.

Following this procedure, we first used the 2MASS extended catalogue to identify galaxies in the IBIS error circle and then we extracted those which were also radio emitters, in order to isolate AGN candidates by means of NVSS and SUMSS radio catalogues. As a result we obtained a set of 8 objects for which we performed a more in-depth study, and in some cases optical and/or X-ray follow up observations, in order to verify their true association with the *INTEGRAL* source as well as their AGN nature and class.

The purpose of this work is not to search out for AGN using different selection criteria, but rather to confirm how a multiwavelength (radio, IR, optical, and X-ray) study of these 8 sources can be used to test the level of reliability and accuracy of the proposed method.

In three cases (IGR J08190–3835, IGR J17520–6018, IGR J21441+4640) we found the X-ray counterparts of the IBIS sources. The optical spectra obtained for two of these sources (IGR J08190–3835, IGR J21441+4640) allowed us to identify them as AGN belonging to the Type 2 and LINER class. The third one (IGR J17520–6018) is most likely a type 2 AGN on the basis of the high X-ray absorption measured. We further suggest that 3 sources (IGR J00556+7708, IGR J17219–1509, IGR J21268+6203) are likely active galaxies on the basis of the radio spectra, NIR photometry and location above the Galactic plane: they are all likely associated with the IBIS objects. LCSB 02890 is instead a starburst galaxy, which is at most a very weak X-ray source and so unlikely to emit at high energies; we conclude that it is an improbable association of IGR J05583–1257. The nature of this *INTEGRAL* source is therefore still open. In only one case (IGR J03103+5706), we have not enough information for a clear classification of the radio/NIR source as an AGN; the nature of this *INTEGRAL* object remains dubious.

Overall, detailed information and follow up measurements confirm the goodness of our method in the search for AGNs among unidentified hard X-ray emitters. On the basis of this work we have classified in the 4th IBIS catalogue the sources discussed in this paper as AGNs and the same approach can be used to pinpoint AGN candidates among Swift/BAT or future *Nustar* (Harrison et al. 2010) unidentified objects.

## ACKNOWLEDGMENTS

We thank José Velásquez for Service Mode observations at the CTIO 1.5m telescope, and Fred Walter for relaying the observing information to him. We also thank the anonymous referee for useful remarks which helped us to improve the quality of this paper. This research has made use of the ASI Science Data Center Multimission Archive; it also used the NASA Astrophysics Data System Abstract Service, the NASA/IPAC Extragalactic Database (NED), and the NASA/IPAC Infrared Science Archive, which are operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. This publication made use of data products from the Two Micron All Sky Survey (2MASS), which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. This research has also made use of data extracted from the Six-degree Field Galaxy Survey and the Sloan Digitized Sky Survey archives; it has also made use of the SIMBAD database operated at CDS, Strasbourg, France, and of the HyperLeda catalog operated at the Observatoire de Lyon, France. The authors acknowledge the ASI and INAF financial support via grant No. I/008/07. LM is supported by the University of Padua through grant No. CPDR061795/06. VC is supported by the CONACYT

research grant 54480-F (México). DM is supported by the Basal CATA PFB 06/09, and FONDAP Center for Astrophysics grant No. 15010003. GG acknowledges the support of Fondecyt regular 1085267, Fondo Gemini 32090009, and Fondo ALMA 31090009. The authors acknowledge the ASI financial support via ASI-INAF contracts I/033/10/0 and I/009/10/0.

Stephen J. B., Bassani L., Landi R., Malizia A., Sguera V., Bazzano A., Masetti N., 2010, MNRAS, 408, 422  
 Ubertini P., et al. 2003, A&A, 411, L131  
 Veilleux S., Osterbrock D.E. 1987, ApJS, 63, 295  
 Winkler C., et al. 2003, A&A, 411, L1  
 Wright E.L. 2006, PASP, 118, 1711

## REFERENCES

- Bird A.J., et al. 2010, ApJS, 186, 1  
 Burrows D. N., et al. 2005, Space Sci. Rev., 120, 165  
 Cardelli J.A., Clayton G.C., Mathis J.S. 1989, ApJ, 345, 245  
 Condon J. J., Cotton W. D., Greisen E. W., Yin Q. F., Perley R. A., Taylor G. B., Broderick J. J., 1998, AJ, 115, 1693  
 Fukugita, M., Shimasaku, K., Ichikawa, T. 1995, PASP, 107, 945  
 Gehrels N., et al. 2004, ApJ, 611, 1005  
 Harrison F. A., et al. 2010, Proc. SPIE, 7732, 21  
 Heckman T.M. 1980, A&A, 87, 152  
 Hill J. E., et al. 2004, Proc. SPIE, 5165, 217  
 Ho L.C., Filippenko A.V., Sargent W.L.W. 1993, ApJ, 417, 63  
 Ho L.C., Filippenko A.V., Sargent W.L.W. 1997, ApJS, 112, 315  
 Horne K. 1986, PASP, 98, 609  
 IRAS, 1988. The Point Source Catalogue, version 2.0, NASA RP-1190  
 Jones D. H., et al. 2004, MNRAS, 355, 747  
 Jones, D. H., Saunders, W., Read, M. Colless, M. 2005, PASA, 22, 277  
 Kalberla P. M. W., Burton W. B., Hartmann Dap, Arnal E. M., Bajaja E., Morras R., Pöppe W. G. L., 2005, A&A, 440, 775  
 Kauffmann G., et al. 2003, MNRAS, 346, 1055  
 Kouzuma S. Yamaoka H. 2010, A&A, 509, A64  
 Lister M. 2003, ASPC, 300, 71  
 Maiorano E., et al. 2010, ATel, 2975, 1  
 Masetti, N., et al. 2010, A&A, 519, A96  
 Mauch T., Murphy T., Buttery H. J., Curran J., Hunstead R. W., Piestrzynski B., Robertson J. G., Sadler E. M., 2003, MNRAS, 342, 1117  
 Mazzarella J.M., Madore B.F., Helou G. and NED team 2001, Proc. SPIE, 4477, 20  
 Monet D.G., et al 2003, AJ, 125, 984  
 Moretti A., et al. 2004, Proc. SPIE, 5165, 232  
 Murakami H., et al. 2007, PASJ, 59S, 369  
 O’Dea, C. 1998, PASP, 110, 493  
 Osterbrock D.E. 1989, Astrophysics of Gaseous Nebulae and Active Galactic Nuclei (Mill Valley: Univ. Science Books)  
 Predehl P., Schmitt J.H.M.M. 1995, A&A, 293, 889  
 Schlegel D.J., Finkbeiner D.P., & Davis M. 1998, ApJ, 500, 525  
 Skrutskie M.F., et al. 2006, AJ, 131, 1163  
 Stanghellini C. 2006, in Proceedings of the 8th VLBI Network Symposium, ed. A. Marecki et al., 18  
 Stephen J. B., et al. 2005, A&A, 432, L49  
 Stephen J. B., et al. 2006, A&A, 445, 869